

Heartworm Preventative & Intestinal Parasite Treatment

QuadriGuard™

(ivermectin/pyrantel pamoate/praziquantel)
Chewable Tablets for Dogs



Exceptional Pet Healthcare

Worm infections cause mild to severe discomfort and can even be fatal – and heartworm is especially dangerous. As well, roundworms, hookworms and tapeworms can be transmitted to humans.

Heartworms: It only takes one bite

Heartworm disease affects dogs throughout the 50 States and can be fatal. Heartworms can start with a bite from a single infected mosquito. Once heartworms mature, they take up residence in the heart and large vessels of the lungs. Heartworms can restrict blood flow throughout the body affecting the heart, lungs, kidneys, and liver.

Signs of Heartworm Disease:

- Coughing
- Lethargy
- Heavy Breathing
- Weight Loss

The good news is heartworm disease is preventable by giving your dog Vethical® QuadriGuard Chewable Tablets for Dogs once a month.



You can feel confident that you are providing the best protection for your dog.

Exceptional Pet Healthcare

The heartworm lifecycle



1. A mosquito bites an infected animal transferring heartworm larvae (microfilariae) into the mosquito as it feeds.

2. The microfilariae develop into infective larvae within the mosquito and are transferred to an uninfected dog when the mosquito bites. Once the infective larvae are in the dog they migrate through the tissue eventually taking residence in the heart.

3. Adult heartworms mature in the heart and begin reproducing microfilariae, starting the cycle again.

All dogs should be tested for heartworm infection before starting a preventive program. Following use of **Vethical QuadriGuard Chewable Tablets for Dogs**, digestive and neurological side effects have rarely been reported. Use with caution in sick, debilitated or underweight animals and dogs weighing less than 10 lbs. See product label for additional information.

Intestinal parasite

Roundworms

Roundworms are the most common intestinal parasite. Your pet can pick up the infection by eating infected soil, licking contaminated fur or paws. Puppies and kittens are the most prone to roundworm infection.

Hookworms

Hookworms are dangerous because they actually bite into the intestinal lining of an animal and suck blood. Left untreated, hookworm infections can result in potentially life-threatening blood loss, weakness, and malnutrition.

CATCH THEM EARLY!


Tapeworms

Tapeworms live in the small intestine and steal the nutrients from your pet's food. Dogs become infected when they eat infected fleas or lice. They can also get certain types of tapeworms by eating infected rodents.

Toxascaris leonina



Toxocara canis



SIGNS Roundworms may be found in your dog's stool or vomit. The symptoms of a severe infestation are: pot belly appearance, diarrhea, vomiting, dull coat and weight loss.

A. braziliense



A. caninum



U. stenocephala



SIGNS A dog infected with hookworm would experience bloody stool, anemia, weight loss, pale gums, diarrhea and a low energy level. Skin irritation can be a sign of a severe infestation.

T. pisiformis



D. caninum



SIGNS Tapeworms can be seen around the dog's anus, in his stool or in his bed. Severe symptoms are abdominal pain, nervousness, severe itching, vomiting and weight loss.

QuadriGuard™

(ivermectin/pyrantel pamoate/praziquantel)
CHEWABLE TABLETS FOR DOGS



For oral use in dogs only.

Caution: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

Description: QuadriGuard™ Chewable Tablets for Dogs is a combination of three anthelmintics (ivermectin/pyrantel pamoate/praziquantel). The tablets are available in four sizes in color-coded packages for oral administration to dogs according to their weight (see **Dosage**).

Indications: For use in dogs to prevent canine heartworm disease by eliminating the tissue stage of heartworm larvae (*Dirofilaria immitis*) for a month (30 days) after infection and for the treatment and control of roundworms (*Toxocara canis*, *Toxascaris leonina*), hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*, *Ancylostoma braziliense*), and tapeworms (*Dipylidium caninum*, *Taenia pisiformis*).

Dosage: QuadriGuard™ (ivermectin/pyrantel pamoate/praziquantel) Chewable Tablets for Dogs should be administered orally at monthly intervals at the recommended minimum dose level of 6 mcg of ivermectin per kilogram (2.72 mcg/lb), 5 mg of pyrantel (as pamoate salt) per kg (2.27 mg/lb) and 5 mg of praziquantel per kg (2.27 mg/lb) of body weight, as follows:

Dog Weight Lbs.	Chewable Tablet Per Month	Tablet Size	Ivermectin Content	Pyrantel Pamoate Content	Praziquantel Content
6.0 to 12	1	Toy	34 mcg	28.5 mg	28.5 mg
12.1 to 25	1	Small	68 mcg	57 mg	57 mg
25.1 to 50	1	Medium	136 mcg	114 mg	114 mg
50.1 to 100	1	Large	272 mcg	228 mg	228 mg

QuadriGuard™ Chewable Tablets for Dogs is recommended for dogs 8 weeks of age and older. For dogs over 100 lbs, use the appropriate combination of these chewable tablets.

Administration: Remove only one tablet at a time from the foil-backed blister card. Return the remaining tablets to their box to protect the product from light. QuadriGuard™ Chewable Tablets for Dogs can be offered to the dog by hand or be added intact to a small amount of dog food. Care should be taken that the dog consumes the complete dose, and the dog should be observed for a few minutes after administration to ensure that part of the dose is not lost or rejected. If it is suspected that any of the dose has been lost, redosing is recommended.

QuadriGuard™ Chewable Tablets for Dogs should be given at monthly intervals during the period of the year when mosquitoes (vectors), potentially carrying infective heartworm larvae, are active. The initial dose must be given within a month (30 days) after the dog's first exposure to mosquitoes. The final dose must be given within a month (30 days) after the dog's last exposure to mosquitoes.

When replacing another heartworm preventive product in a heartworm disease prevention program, the first dose of QuadriGuard™ Chewable Tablets for Dogs must be given within a month (30 days) of the last dose of the former medication. A heartworm test should be performed prior to switching heartworm preventive products.

If the interval between doses exceeds a month (30 days), the efficacy of ivermectin can be reduced. Therefore, for optimal performance, the tablet must be given once a month on or about the same day of the month. If treatment is delayed, whether by a few days or many, immediate treatment with QuadriGuard™ Chewable Tablets for Dogs and resumption of the recommended dosing regimen will minimize the opportunity for the development of adult heartworms.

WARNING:

For use in dogs only. Keep this and all drugs out of reach of children. In safety studies, testicular hypoplasia was observed in some dogs receiving 3 and 5 times the maximum recommended dose monthly for 6 months (see Animal Safety).

In case of ingestion by humans, clients should be advised to contact a physician immediately. Physicians may contact a Poison Control Center for advice concerning cases of ingestion by humans.

For a copy of the Material Safety Data Sheet (MSDS), or to report adverse reactions, call 1-866-337-5009.

Precautions: Use with caution in sick, debilitated, or underweight animals and dogs weighing less than 10 lbs (see Animal Safety). The safe use of this drug has not been evaluated in pregnant or lactating bitches.

All dogs should be tested for existing heartworm infection before starting treatment with QuadriGuard™ Chewable Tablets for Dogs, which is not effective against adult *D. immitis*. Infected dogs should be treated to remove adult heartworms and microfilariae before initiating a heartworm prevention program.

While some microfilariae may be killed by the ivermectin in QuadriGuard™ Chewable Tablets for Dogs at the recommended dose level, QuadriGuard™ Chewable Tablets for Dogs is not effective for microfilariae clearance. A mild hypersensitivity-type reaction, presumably due to dead or dying microfilariae and particularly involving a transient diarrhea, has been observed in clinical trials with ivermectin alone after treatment of some dogs that have circulating microfilariae.

Adverse Reactions: Self-limiting adverse reactions including lethargy, limpness, salivation, shaking, diarrhea, decreased appetite, licking lips, and belching were reported between 20 minutes and 72 hours following treatment in a field study with QuadriGuard™ Chewable Tablets for Dogs.

In clinical field trials with ivermectin/pyrantel pamoate, vomiting or diarrhea within 24 hours of dosing was rarely observed (1.1% of administered doses). The following adverse reactions have been reported following the use of ivermectin: depression/lethargy, vomiting, anorexia, diarrhea, mydriasis, ataxia, staggering, convulsions and hypersalivation.

Effectiveness: QuadriGuard™ Chewable Tablets for Dogs, given orally using the recommended dose and regimen, is effective against the tissue larval stage of *D. immitis* for a month (30 days) after infection and, as a result, prevent the development of the adult stage. QuadriGuard™ Chewable Tablets for Dogs is also effective against canine roundworms (*T. canis*, *T. leonina*), hookworms (*A. caninum*, *U. stenocephala*, *A. braziliense*), and tapeworms (*T. pisiformis*, *D. caninum*). A total of 61 dogs and puppies with naturally acquired or experimental parasite infections treated with QuadriGuard™ Chewable Tablets for Dogs were enrolled in 7 well-controlled laboratory studies to establish effectiveness. These studies included dogs of various sizes, ages and breeds. Data from these studies demonstrated QuadriGuard™ Chewable Tablets for Dogs is safe and effective for the removal of the parasite species indicated on the label when used as directed.

Palatability: In a field trial, QuadriGuard™ was shown to be a palatable oral dosage form for dogs.

Animal Safety: Studies with ivermectin indicate that certain dogs of the Collie breed are more sensitive to the effects of ivermectin administered at elevated dose levels (more than 16 times the target use level of 6 mcg/kg) than dogs of other breeds. At elevated doses, sensitive dogs showed adverse reactions which included mydriasis, depression, ataxia, tremors, drooling, paresis, recumbency, excitability, stupor, coma and death. No signs of toxicity were seen at 10 times the recommended dose (27.2 mcg/lb) in sensitive Collies. Results of these studies and bioequivalence studies support the safety of ivermectin products in dogs, including Collies, when used as recommended by the label.

In a target animal safety study, QuadriGuard™ Chewable Tablets for Dogs was administered to 8-week-old Beagle puppies at one, three and five times the maximum recommended dose of 12.5 mcg/kg ivermectin, 10.47 mg/kg pyrantel and 10.47 mg/kg praziquantel. The dogs were treated every 30 days for 6 months. Vomiting within 6 hours of dosing and soft or watery feces within 24 hours of dosing were observed. Other observations during the study were: anogenital swelling, lethargy, head movements, shallow, audible or difficult breathing, and salivation. One dog in the 5X group had tremors and decreased activity. All these signs were transient. No treatment was required. Pathology showed testicular hypoplasia in the 3 and 5X groups (see Warnings).

In a laboratory safety study, 12-week-old Beagle puppies receiving 3 and 5 times the recommended dose once weekly for 13 weeks demonstrated a dose-related decrease in testicular maturation compared to controls. In this study, all treated puppies had significantly higher cholesterol levels compared to untreated controls.

In a reproductive safety study, adult males were treated at 37.5 mcg/kg ivermectin, 31.4 mg/kg pyrantel and 31.4 mg/kg praziquantel every 14 days during two full spermatogenic cycles (112 days). The quality of semen and reproductive health were not affected by treatment. Treatment-related vomiting and soft feces were reported during this study.

In a study of the effectiveness of QuadriGuard™ Chewable Tablets for Dogs for treatment of *T. canis*, one 8.1 lb, 72-day-old puppy died 6 days after administration of QuadriGuard™ Chewable Tablets for Dogs at the label dose. This puppy and many of the other puppies in the study had high worm burdens and were reported to have diarrhea, sometimes bloody, frequently before and after treatment with QuadriGuard™ Chewable Tablets for Dogs or the placebo. Dehydration and signs of anemia (pale mucous membranes) were the only abnormal gross necropsy findings observed. No definitive cause of death was determined. In a 90-day field study, the most serious ADEs (lethargy, limpness, and salivation) were seen in dogs weighing less than 10 lbs (see Precautions).

Storage Information: Store at 20°C-25°C (68°F-77°F), excursions permitted between 15°C-30°C (59°F-86°F). Protect product from light.

How Supplied: QuadriGuard™ Chewable Tablets for Dogs is available in four dosage strengths (see dosage section) for dogs of different weights. Each strength comes in a box of 6 chewable tablets, packed 10 boxes per display box.

Make sure your dog is protected, with the guidance of your veterinarian and with Vethical QuadriGuard Chewable Tablets for Dogs – a top-quality, once-a-month tablet you can count on.



- **Broad-spectrum** - The most complete protection available against heartworm, two species of roundworm, three species of hookworm, and two species of tapeworm (one of the only products available on the market to offer this level of protection).
- **Easy to give** - Dogs love the flavored chewable tablets — and it is safe to treat puppies even as young as eight weeks.
- **Proven and safe** - Guaranteed protection and satisfaction when used as directed.
- **FDA-approved**

VCA Home Delivery Simple!

Parasite Protection

- Single-dose, monthly delivery
- Pay as you go
- You don't pay more than in the hospital
- No more stickers, calendars, or emails, VCA Home Delivery is your best reminder
- **FREE shipping***
- **PLUS get a FREE semi-annual health exam****

**FREE
shipping*, plus
Semi-Annual
Exam****

* Ask for details.

** Valid with purchase of Flea, Tick, and/or Heartworm monthly delivery.
Some conditions apply. Ask for details.

For more information, go to **VETHICAL.COM**

3010-PL-FL

Exceptional Pet Healthcare